

## A review on phytochemistry and pharmacological effects of *Trigonella foenum-graecum*

Ayah Rebhi Hilles<sup>1\*</sup>, Syed Mahmood<sup>2</sup>

<sup>1</sup>Physiology Dept., Kulliyyah of Medicine, International Islamic University Malaysia (IIUM), Kuantan, Pahang, Malaysia; <sup>2</sup>Pharmaceutical Technology Dept., Kulliyyah of Pharmacy, International Islamic University Malaysia, Kuantan, Pahang, Malaysia.

Received: 2/Aug/2016 Accepted: 21/Sep/2016

### ABSTRACT

**Background and aims:** Fenugreek (*Trigonella foenum-graecum*) is a medicinal plant with potential applications in the medicinal purposes. Fenugreek has been used in food as a flavouring agent since ancient times in many countries like Greece, Rome and Egypt. It has a special place in the traditional medical system. The aim of this study was to review phytochemistry and pharmacological effects of *Trigonella foenum-graecum*.

**Methods:** This research was a review study. Data were collected using articles in various databases and books about pharmacological effects of *Trigonella foenum-graecum*.

**Results:** The seeds are rich sources of lipids, protein, mucilage, calcium, dietary fiber B vitamins, Iron, protease inhibitors and several steroid saponins, tiny amounts of alkaloid, furostanol glycosides and steroidal peptide. As a medicinal plant, fenugreek recognized as a galactagogue or lactation stimulant in women after child birth as well as for its ability to treat wounds and sore muscles. In addition, it has the ability to be as antibacterial, anti-ulcer, anti-cancer, anthelmintic (antagonistic effect against parasitic worms), and antinociceptive (pain-reducing) properties. In recent years, laboratory studies and clinical trials have focused on the potential activity of fenugreek as a natural medicine.

**Conclusion:** These studies have shown that fenugreek plants possess immunomodulatory, hypocholesterolaemic, hypoglycaemic, gastro- and hepatoprotective and antioxidative properties. Pharmacological properties of fenugreek have been explored to identify a role for the plant in diabetes management and in cardiovascular health, indicating the presence of bioactive compounds in fenugreek, which may be responsible for its health benefits.

**Keywords:** Fenugreek, Galactagogue, Antinociceptive, Hepatoprotective, Hypoglycaemic, Antioxidative.

### INTRODUCTION

Fenugreek (*Trigonella foenum-graecum* L.) is annual plant from Fabaceae family, which is native to the Indian subcontinent and the Eastern Mediterranean region.<sup>1</sup> Fenugreek,

is known for presence of the distinctive aromatic compounds that gives special flavour and colour to the food.<sup>2</sup> Fresh fenugreek leaves consider as an ingredient in

\*Corresponding author: Ayah Rebhi Hilles. Physiology Dept., Kulliyyah of Medicine, International Islamic University Malaysia (IIUM), Kuantan, Pahang, Malaysia, Tel: 0060178322697, E-mail: ayah.hilles90@gmail.com

some Indian Currie.<sup>3</sup> It is one of the most ancient medicinal herbs. It provides natural food fibre and other nutrients required in human body.<sup>4</sup>

Fenugreek is one of the oldest known medicinal plants which has been documented in ancient herbal history.<sup>1</sup> Seeds of the fenugreek have been used as a holy smoke that Egyptians consumed in their embalming rites during Pharaohs time.<sup>5</sup> It has been used also to promote labour before delivery during Greek period.<sup>6</sup> According to Chinese traditional medicine, fenugreek can be used to treat Lymphedema (oedema of the legs).<sup>7</sup>

Fenugreek seed is a good source of calcium, minerals, iron,  $\beta$ -carotene and several vitamins like vitamins A and D. It is rich source of available carbohydrates and dietary fiber.<sup>8</sup> It is a source of free amino acids; 4-hydroxyisoleucine, lysine, histidine and arginine (25.8%), protein (20-30%), moisture (11.76%), fat (6.53%), crude fibre (6.28%), ash content (3.26%) and energy (394.46 Kcal/100 g seed).<sup>9</sup> It contains lecithin, choline, minerals, B. Complex, Phosphates, and Para-Amino Benzoic acid (PABA). In addition, the main chemical compounds in fenugreek are saponins, fenugreekine, trigonelline, coumarin, scopoletin, phytic acid and nicotinic acid.<sup>10</sup> The significance of *T. foenum-graecum* seeds is due to the defatted part, with high quality fibre including steroidal saponins and protein comparable to those of soybean.<sup>11</sup> The important chemical constituents are saponins, coumarin, fenugreekine, nicotinic acid, phytic acid, scopoletin and trigonelline. The seeds also have the alkaloid trigonelline with mucilage, tannic acid, yellow colour substance, fixed and volatile oils and a bitter extractive, diosgenin and gitogenin a trace of trigogenin.<sup>12</sup> The main bioactive compounds in fenugreek are Galactomannan, Diosgenin, 4-Hydroxyisoleucine.<sup>13</sup>

Fenugreek showed a mild hypoglycemic effect as well as improvement in glucose homeostasis and reduction in low-density lipoprotein (LDL), very-low-density lipoprotein (VLDL), high-density lipoprotein (HDL) and total cholesterol in alloxan diabetic rats.<sup>14-16</sup> It can be used as a control for diabetes mellitus type 2.<sup>17</sup> It has been proven that the soluble dietary fiber of *fenugreek seeds* has antidiabetic effect mediated through enhancement of peripheral insulin action and inhibition of carbohydrate digestion and absorption.<sup>18</sup> The antidiabetic effect of fenugreek has been recorded as same as to glibenclamide treatment which is antidiabetic drug, this shows that fenugreek can be considered as a candidate for future studies on diabetes mellitus.<sup>19</sup>

It has been reported that fenugreek has antioxidant activity.<sup>20,21</sup> It also can be considered as a potent antioxidant.<sup>22</sup> Due to normalize the disruption in free radical metabolism in diabetic rats, and the presence of flavonoids; especially malondialdehyde, nonprotein thiol, protein, and cysteine contents in the plant, it has been shown that fenugreek seeds contain antioxidants which protect cellular structures from oxidative damage and alterations in antioxidant enzymes.<sup>23-28</sup> Fenugreek exerts its chemopreventive effect by decreasing circulatory lipoproteins and enhancing antioxidant levels.<sup>29</sup>

It has been shown the hypolipidaemic effects of fenugreek seeds due to the fraction rich in fibres (hypocholesterolaemic and hypotriglyceridaemic effects), and to the fraction rich in saponins.<sup>30,31</sup> It has been proven that fenugreek increases the plasma insulin, decreases the total cholesterol, very low-density lipoprotein (VLDL) and low-density lipoprotein (LDL).<sup>32,33</sup> Also, some other various medicinal and pharmacologic properties of fenugreek which have been stated but received less attention including anti-bacterial,

anti-cholinergic, anthelmintic, wound healing activities.<sup>34</sup>

In recent years, various dietary components that can potentially be used for the prevention and treatment of cancer have been identified.<sup>35</sup> It has been proved that diosgenin induced apoptosis in HT-29 cells at least in part by inhibition of bcl-2 and by induction of caspase-3 protein expression. On the basis of these findings, the fenugreek constituent diosgenin seems to have potential as a novel colon cancer preventive agent.<sup>36</sup>

Greek showed cytotoxic activity against various cancer cell lines.<sup>37</sup> It killed MCF-7 human breast cancer cells via an apoptotic pathway.<sup>38,39</sup> Protodioscin (PD) was purified from fenugreek (*Trigonella foenum-graecum* L.) and identified by Mass, and 1H- and 13C-NMR. The effects of PD on cell viability in human leukemia HL-60 and human stomach cancer KATO III cells were investigated. PD displayed strong growth inhibitory effect against HL-60 cells, these findings suggest that growth inhibition by PD of HL-60 cells results from the induction of apoptosis by this compound in HL-60 cells Fenugreek has anti-Leukemic potential activity.<sup>40,41</sup>

It has been demonstrated that fenugreek has anti-inflammatory and antipyretic properties.<sup>42</sup> As it can protect against skin damage because it is strongly suppressed the production of inflammatory cytokines.<sup>43</sup> Treated arthritic rats with fenugreek showed an increase in ESR and total WBC, a decrease in RBC count and hemoglobin and aberrant changes to the C-reactive protein levels.<sup>44</sup> The outcoming results of phytochemical tests confirmed the presence of flavonoids in fenugreek seeds which is the responsible for anti-inflammatory activity of the plant.<sup>45</sup> Fenugreek seeds have antioxidant, anti-inflammatory, and antineoplastic activities.<sup>46-48</sup> The recent studies support the traditional uses of fenugreek for inflammations. However, more researchs are needed for its use in clinical studies.<sup>49</sup>

It has been reported that fenugreek has antiandrogenic and antifertility potential activities.<sup>50,51</sup> It has been documented that fenugreek seeds have antifertility effect in the female rabbits and more of a toxicity effect in the male rabbits.<sup>52</sup> The antifertility activity of fenugreek exists because of the presence of saponins.<sup>53</sup> It has been showed that fenugreek seeds are effective almost three times more than combined oral contraceptive pills at higher dose.<sup>54</sup>

A 30% of fenugreek powder for 20 days lead to reduction in the fetal and placental weights.<sup>55</sup> A 3.2 g/kg dosage of fenugreek affects on the bone marrow proliferation; bone marrow cells have not been clearly demonstrated.<sup>56</sup> A 0.11 g/kg of fenugreek for 15 days to male rats could lead to the changes in the level of thyroid hormones.<sup>57</sup> Fenugreek aqueous extract affects the fertility by promoting anti-implantation, or through embryonic loss or re-absorption.<sup>58,59</sup> At the 2 doses (500 and 1000 mg/kg/day) of fenugreek lyophilized aqueous extract increased pup mortality and reduced body weights.<sup>60</sup>

## CONCLUSION

Fenugreek is an annual plant in the family Fabaceae. Fenugreek has a long history as medicinal herb in the ancient world. The largest producer of fenugreek seeds is India. Its seeds are a common ingredient in dishes from the Indian subcontinent. The most three bioactive compounds that were examined in experiments are: Galactomannan, diosgenin and 4-hydroxyisoleucine. Fenugreek has potential therapeutic activities in so many treatments such as diabetes, atherosclerosis, digestive problems, chronic coughs, inflammation, milk flow in Breastfeeding, tuberculosis, baldness, Parkinson's disease, cancer, and exercise performance.

## REFERENCES

1. Petropoulos GA. Fenugreek, the Genus *Trigonella*, Medicinal and Aromatic Plants-Industrial Profiles. TJ International Ltd., Padstow, Cornwall. 2002.
2. Max B. This and that: The essential pharmacology of herbs and spices. Trends Pharmacol Sci. 1992; 13:15-20.
3. Srinivasan K. Fenugreek (*Trigonella foenum-graecum*): A review of health beneficial physiological effects. Food Rev Int. 2006 Jul 1; 22(2): 203-24.
4. Thomas JE, Bandara M, Lee EL, Driedger D, Acharya S. Biochemical monitoring in fenugreek to develop functional food and medicinal plant variants. N Biotechnol. 2011; 28(2): 110-7.
5. Fazli F, Hardman R. The spice, fenugreek (*Trigonella foenum-graecum* L.): Its commercial varieties of seed as a source mehfafarin & authors of diosgenin. Trop Sci. 1968; 10: 66 - 78.
6. Yoshikawa M, Murakami T, Komatsu H, Murakami N, Yamahara J, Matsuda H. Medicinal foodstuffs. IV. Fenugreek seed. (1): Structures of trigoneosides Ia, Ib, IIa, IIb, IIIa, and IIIb, new furostanol saponins from the seeds of Indian *Trigonella foenum-graecum* L. Chem Pharm Bull (Tokyo). 1997; 45(1): 81-7.
7. Smith M. Therapeutic applications of fenugreek. Alt Med Rev. 2003; 8(1): 20-7.
8. Kochhar A, Nagi M, Sachdeva R. Proximate composition, available carbohydrates, dietary fibre and anti nutritional factors of selected traditional medicinal plants. J Hum Ecol. 2006; 19(3): 195-9.
9. Khan FU, Ullah A, Rahman S, Naz S, Rana N. Fenugreek (*Trigonella foenum-graecum* L.) effect on muscle growth of broiler chicks. Res Opin Anim Vet Sci. 2011. Available from: <http://scholar.google.com>.
10. Valette G, Sauvaire Y, Baccou JC, Ribes G. Hypocholesterolaemic effect of fenugreek seeds in dogs. Atherosclerosis. 1984; 50(1): 105-11.
11. Varel VH, Nienaber JA, Freetly HC. Conservation of nitrogen in cattle feedlot waste with urease inhibitors. J Anim Sci. 1999; 77(5): 1162-8.
12. Jayaweera DM. Medicinal Plants (Indigenous and Exotic) used in Ceylon. Colombo: The National Science Council of Sri Lanka; 1981.
13. Lee EL. Genotype X environment impact on selected bioactive compound content of fenugreek (*Trigonella foenum-graecum* L.) [Dissertation]. Lethbridge, Alta.: University of Lethbridge, Dept. of Biological Sciences; 2009.
14. Jelodar GA, Maleki M, Motadayen MH, Sirus S. Effect of fenugreek, onion and garlic on blood glucose and histopathology of pancreas of alloxan-induced diabetic rats. Indian J Med Sci. 2005; 59(2): 64-9.
15. Raju J, Gupta D, Rao AR, Yadava PK, Baquer NZ. *Trigonella foenum graecum* (fenugreek) seed powder improves glucose homeostasis in alloxan diabetic rat tissues by reversing the altered glycolytic, gluconeogenic and lipogenic enzymes. Mol Cell Biochem. 2001; 224(1-2): 45-51.
16. Al-Habori M, Raman A. Antidiabetic and hypocholesterolaemic effects of fenugreek. Phytother Res. 1998; 12(4): 233-42.
17. Kassaian N, Azadbakht L, Forghani B, Amini M. Effect of fenugreek seeds on blood glucose and lipid profiles in type 2 diabetic patients. Int J Vitam Nutr Res. 2009; 79(1): 34-9.
18. Hannan JM, Ali L, Rokeya B, Khaleque J, Akhter M, Flatt PR, et al. Soluble dietary fibre fraction of *Trigonella foenum-graecum* (fenugreek) seed improves glucose homeostasis in animal models of type 1 and type 2 diabetes by delaying carbohydrate digestion and absorption, and enhancing insulin action. Br J Nutr. 2007; 97(3): 514-21.
19. Eidi A, Eidi M, Sokhteh M. Effect of fenugreek (*Trigonella foenum-graecum* L.) seeds on serum parameters in normal and

- streptozotocin-induced diabetic rats. *Nutr Res.* 2007; 27(11): 728-733.
20. Rababah TM, Hettiarachchy NS, Horax R. Total phenolics and antioxidant activities of fenugreek, green tea, black tea, grape seed, ginger, rosemary, gotu kola, and ginkgo extracts, vitamin E, and tert-butylhydroquinone. *J Agric Food Chem.* 2004; 52(16): 5183-6.
21. Anuradha CV, Ravikumar P. Restoration on tissue antioxidants by fenugreek seeds (*Trigonella Foenum Graecum*) in alloxan-diabetic rats. *Indian J Physiol Pharmacol.* 2001; 45(4): 408-20.
22. Bukhari SB, Bhangar MI, Memon S. Antioxidative activity of extracts from fenugreek. *Pak J Anal Environ Chem.* 2008; 9: 78-83.
23. Ravikumar P, Anuradha CV. Effect of fenugreek seeds on blood lipid peroxidation and antioxidants in diabetic rats. *Phytother Res.* 1999; 13(3): 197-201.
24. Randhir R, Lin YT, Shetty K. Phenolics, their antioxidant and antimicrobial activity in dark germinated fenugreek sprouts in response to peptide and phytochemical elicitors. *Asia Pac J Clin Nutr.* 2004. 13(3): 295-307.
25. Sinha S, Gupta AK, Bhatt K. Uptake and translocation of metals in fenugreek grown on soil amended with tannery sludge: involvement of antioxidants. *Ecotoxicol Environ Saf.* 2007; 67(2): 267-77.
26. Naidu MM, Shyamala BN, Naik JP, Sulochanamma G, Srinivas P. Chemical composition and antioxidant activity of the husk and endosperm of fenugreek seeds. *LWT-Food Sci Technol.* 2011; 44(2): 451-6.
27. Kaviarasan S, Naik GH, Gangabhairathi R, Anuradha CV, Priyadarsini KI. *In vitro* studies on antiradical and antioxidant activities of fenugreek (*Trigonella foenum graecum*) seeds. *Food chem.* 2007; 103(1): 31-7.
28. Genet S, Kale RK, Baquer NZ. Alterations in antioxidant enzymes and oxidative damage in experimental diabetic rat tissues: effect of vanadate and fenugreek (*Trigonella foenum graecum*). *Molecul cell Biochem.* 2002; 236(1-2): 7-12.
29. Devasena T, Menon VP. Enhancement of circulatory antioxidants by fenugreek during 1, 2-dimethylhydrazine-induced rat colon carcinogenesis. *Journal of biochemistry, molecular biology, and biophysics: J Biochem Mol Biol Biophys.* 2002; 6(4): 289-92.
30. Ribes G, Da Costa C, Loubatières-Mariani MM, Sauvaire Y, Baccou JC. Hypocholesterolaemic and hypotriglyceridaemic effects of subfractions from fenugreek seeds in alloxan diabetic dogs. *Phytother Res.* 1987; 1(1): 38-43.
31. Petit PR, Sauvaire YD, Hillaire-Buys DM, Leconte OM, Baissac YG, Ponsin GR, et al. Steroid saponins from fenugreek seeds: Extraction, purification, and pharmacological investigation on feeding behavior and plasma cholesterol. *Steroids.* 1995; 60(10): 674-80.
32. Petit P, Sauvaire Y, Ponsin G, Manteghetti M, Fave A, Ribes G. Effects of a fenugreek seed extract on feeding behaviour in the rat: Metabolic-endocrine correlates. *Pharmacol Biochem Behav.* 1993; 45(2): 369-74.
33. Al-Habori M, Al-Aghbari AM, Al-Mamary M. Effects of fenugreek seeds and its extracts on plasma lipid profile: A study on rabbits. *Phytother Res.* 1998; 12(8): 572-5.
34. Adedapo AA, Ofuegbe SO, Soetan KO. Pharmacologic and medicinal properties of fenugreek (*Trigonella Foenum-Graecum* L.). *Am J Soc Issues Human.* 2014.
35. Shabbeer S, Sobolewski M, Anchoori RK, Kachhap S, Hidalgo M, Jimeno A, et al. Fenugreek: A naturally occurring edible spice as an anticancer agent. *Cancer Biol Therap.* 2009; 8(3): 272-8.
36. Raju J, Patlolla JM, Swamy MV, Rao CV. Diosgenin, a steroid saponin of *Trigonella foenum graecum* (Fenugreek), inhibits azoxymethane-induced aberrant crypt foci formation in F344 rats and induces apoptosis in HT-29 human colon

cancer cells. *Cancer Epidemiol Biomarkers Prev.* 2004; 13(8): 1392-8.

37. Al-Oqail MM, Farshori NN, Al-Sheddi ES, Musarrat J, Al-Khedhairi AA, Siddiqui MA. *In vitro* cytotoxic activity of seed oil of fenugreek against various cancer cell lines. *Asian Pac J Cancer Prev.* 2013; 14(3): 1829-32.

38. Sebastian KS, Thampan RV. Differential effects of soybean and fenugreek extracts on the growth of MCF-7 cells. *Chem Biol Interact.* 2007; 170(2): 135-43.

39. Khoja KK, Shaf G, Hasan TN, Syed NA, Al-Khalifa AS, Al-Assaf AH, et al. Fenugreek, a naturally occurring edible spice, kills MCF-7 human breast cancer cells via an apoptotic pathway. *Asian Pac J Cancer Prev.* 2011; 12(12): 3299-304.

40. Hiroshigehibasami HM, Ishikawa K, Katsuzaki H, Kunioimai K, Ishii Y, YA TK. Protodioscin isolated from fenugreek *Trigonella foenum graecum* induces cell death and morphological change indicative of apoptosis in leukemic cell line H-60, but not in gastric cancer cell line KATO III. *Int J Molecul Med.* 2003; 11: 23-6.

41. Gopal PK, Paul M, Paul S. Anti Leukemic potential of different fenugreek seed germplasms. *Am J Soc Issues Human.* 2014.

42. Ahmadiani A, Javan M, Semnianian S, Barat E, Kamalinejad M. Anti-inflammatory and antipyretic effects of *Trigonella foenum-graecum* leaves extract in the rat. *J Ethnopharmacol.* 2001; 75(2): 283-6.

43. Kawabata T, Cui MY, Hasegawa T, Takano F, Ohta T. Anti-inflammatory and anti-melanogenic steroidal saponin glycosides from Fenugreek (*Trigonella foenum-graecum* L.) seeds. *Planta medica.* 2011; 77(7): 705-10.

44. Sindhu G, Ratheesh M, Shyni GL, Nambisan B, Helen A. Anti-inflammatory and antioxidative effects of mucilage of *Trigonella foenum graecum* (Fenugreek) on adjuvant induced arthritic rats. *Int immunopharmacol.* 2012; 12(1): 205-11.

45. Mandegary A, Pournamdari M, Sharififar F, Pournourmohammadi S, Fardiar R, Shooli S. Alkaloid and flavonoid rich fractions of fenugreek seeds (*Trigonella foenum-graecum* L.) with antinociceptive and anti-inflammatory effects. *Food and chem toxicol.* 2012; 50(7): 2503-7.

46. Liu Y, Kakani R, Nair MG. Compounds in functional food fenugreek spice exhibit anti-inflammatory and antioxidant activities. *Food Chem.* 2012 Apr 15; 131(4): 1187-92.

47. Vyas S, Agrawal RP, Solanki P, Trivedi P. Analgesic and anti-inflammatory activities of *Trigonella foenum-graecum* (seed) extract. *Acta Pol Pharm.* 2008; 65(4): 473-6.

48. Sur P, Das M, Gomes A, Vedasiromoni JR, Sahu NP, Banerjee S, et al. Ganguly DK. *Trigonella foenum graecum* (fenugreek) seed extract as an antineoplastic agent. *Phytother Res.* 2001; 15(3): 257-9.

49. Sharififar F, Khazaeli P, Alli N. *In vivo* evaluation of anti-inflammatory activity of topical preparations from Fenugreek (*Trigonella foenum-graecum* L.) seeds in a cream base. *Iranian J Pharmaceut Sci.* 2009; 5(3): 157-62.

50. Kamal R, Yadav R, Sharma JD. Efficacy of the steroidal fraction of fenugreek seed extract on fertility of male albino rats. *Phytother Res.* 1993; 7(2): 134-8.

51. Joshi SC, Sharma A, Chaturvedi M. Antifertility potential of some medicinal plants in males: An overview. *Int J Pharm Pharmaceu Sci.* 2011; 3: 204-17.

52. Kassem A, Al-Aghbari A, M AL-H, Al-Mamary M. Evaluation of the potential antifertility effect of fenugreek seeds in male and female rabbits. *Contraception.* 2006; 73(3): 301-6.

53. Dande PA, Patil SU. Evaluation of Saponins from *Trigonella foenum-graecum* Seeds for its Antifertility Activity. *Asian J Pharm Clin Res.* 2012; 5(3): 154-7.

54. Hilles AR, Allow AK, Mahmood S. Evaluation and comparison of the antifertility potential activity and adverse

effects of *Trigonella foenum-graecum* seeds and combined oral contraceptive pills in female rats. Int J Reprod Contracept Obstet Gynecol. 2016; 5(3): 680-8.

55. Kassem A, Al-Aghbari A, M AL-H, Al-Mamary M. Evaluation of the potential antifertility effect of fenugreek seeds in male and female rabbits. Contraception. 2006; 73(3): 301-6.

56. Araee M, Norouzi M, Habibi G, Sheikhvatan M. Toxicity of *Trigonella foenum-graecum* (Fenugreek) in bone marrow cell proliferation in rat. Pak J Pharm Sci. 2009; 22(2): 126-30.

57. Kelly GS. Peripheral metabolism of thyroid hormones: A review. Altern Med Rev. 2000; 5(4): 306-33.

58. Chamorro G, Salazar M, Fournier G, Pages N. The anti-implantation effects of various savine extracts on the pregnant rat. J Toxicol Clin Exp. 1990; 10(3): 157-60.

59. Asongalem EA, Nana P, Foyet HS, Dimo T, Kamtchouing P. Antifertility and fetotoxic activities of *Acanthus montanus* aqueous extract in Wistar rats. Methods Find Exp Clin Pharmacol. 2008; 30(7): 521-8.

60. Khalki L, M'Hamed S B, Bennis M, Chait A, Sokar Z. Evaluation of the developmental toxicity of the aqueous extract from *Trigonella foenum-graecum* (L.) in mice. J Ethnopharmacol. 2010; 131(2): 321-5.

**How to cite the article:** Hilles AR, Mahmood S. A review on phytochemistry and pharmacological effects of *Trigonella foenum-graecum*. Adv Herb Med. 2017; 3(3): 61-67.